

Probabilistic properties of nonextensive thermodynamic

Franck Jedrzejewski

Commissariat à l'énergie atomique
CEA Saclay, INSTN, F-91191 Gif-sur-Yvette, France
franck@arthaud.saclay.cea.fr

Abstract

Based on the Tsallis entropy, the nonextensive thermodynamic properties are studied as a q -deformation of classical statistical results using only probabilistic methods and straightforward calculations. It is shown that the constant in the Tsallis entropy depends on the deformation parameter and must be redefined to recover the usual thermodynamic relations. The notions of variance and covariance are generalised. A partial derivative formula of the entropy is established. It verifies important relations from which most of the nonextensive thermodynamics relations can be recovered. This leads to a new proof of a highly nontrivial conclusion that thermodynamic relations and Maxwell relations in non extensive thermodynamic have the same forms as those in ordinary nonextensive statistical mechanics. Theoretical results are applied to ideal gas. The case of fermions and bosons systems with fractal distributions is also considered.

PACS : 05.20.-y, 05.70.-a, 05.30.-d.

1 Introduction

The purpose of this work is to discuss the thermodynamic relations problem in Tsallis thermostatics recently presented in Ref. [1]. It has been pointed out that q -thermostatics could be understood in terms of q -deformations. The Tsallis' entropy proposed by Constantino Tsallis ten years ago [2] has found many applications in various domains of physics [3], as the turbulence in electron plasmas, phonon-electron thermalization in ion bombarded solids and flux of solar neutrinos. There are also applications to the mechanisms of anomalous diffusion [4]. When sys-

tems involve long-range interactions, statistical mechanics presents serious difficulties due to non-markovianity of stochastic processes. Non-extensive thermodynamics may correctly cover some of the known anomalies. The canonical ensemble of non-extensive thermodynamics for quantum mechanical systems with finite number of degrees of freedom has been studied in Ref. [5]. It has been shown that most of thermodynamic relations are preserved [6], if introducing the entropic index q , we replace the usual means by q -expectations. However, invariant relations involve Boltzmann-Gibbs constant in the canonical ensemble and are related to Tsallis constant in the grand canonical ensemble. The canonical distribution in non-extensive thermodynamic is given by optimal Lagrange multipliers and is discussed in Ref. [7]. Parameter differentiation of fractional powers of operators has been developed in Ref. [8, 9]. Nonextensive statistical mechanics and thermodynamics, as a new stream in the research of the foundations of statistical mechanics, is presented in a recent publication [10].

2 Tsallis Entropy

The Tsallis' entropy is defined by the formula

$$S_q = k \int \rho \ln_q(\rho^{-1}) d\lambda \quad (1)$$

where ρ is the distribution of probability, λ is the Lebesgue measure on the phase space and k is a constant, called the Tsallis constant. In this expression the usual logarithmic function is replaced by the q -logarithmic function

$$\ln_q(x) = \frac{x^{1-q} - 1}{1 - q} \quad (2)$$

which is the inverse function of the q -exponential function

$$e_q(x) = (1 + (1 - q)x)^{1/(1-q)} \quad (3)$$

defined if $(1 + (1 - q)x) > 0$ and which is prolonged by continuity with $e_q(x) = 0$ otherwise. When $q \rightarrow 1$, the q -exponential and the q -logarithmic functions converge to the ordinary exponential and logarithmic functions. The entropy of Tsallis reduces to the Boltzmann-Gibbs formula in the extensive limit. The pseudo-additivity of the q -logarithmic function

$$\ln_q(xy) = \ln_q(x) + \ln_q(y) + (1 - q) \ln_q(x) \ln_q(y) \quad (4)$$

leads to the nonextensivity of the Tsallis entropy. For two independent subsystems A and B of a total system, in thermal contact with each other, it has been shown that

$$S_q(\rho_A \otimes \rho_B) = S_q(\rho_A) + S_q(\rho_B) + (1 - q)k S_q(\rho_A) S_q(\rho_B) \quad (5)$$

The following property of the q -logarithmic function

$$\ln_q\left(\frac{1}{x}\right) = -x^{q-1} \ln_q(x) \quad (6)$$

gives the usual definition of the Tsallis entropy [11]

$$S_q = -k \int \rho^q \ln_q(\rho) d\lambda \quad (7)$$

which is also defined by the equivalent formula

$$S_q = \frac{k}{q-1} \left(1 - \int \rho^q d\lambda \right) \quad (8)$$

If we define the unnormalized q -expectation of an observable A by

$$[A]_q = \int A \rho^q d\lambda \quad (9)$$

and let c_q be the constant

$$c_q = \int \rho^q d\lambda \quad (10)$$

the q -expectation (normalised) could be defined by

$$\langle A \rangle_q = \frac{[A]_q}{c_q} = \frac{1}{c_q} \int A \rho^q d\lambda \quad (11)$$

The problem posed by unnormalized expectation is studied in Ref. [12]. Introducing the q -Gibbs measure (Ref. [13]) in the canonical ensemble

$$\rho = \frac{e_q(-\beta H)}{Z_q} \quad (12)$$

where H is the Hamiltonian of the system and β the Lagrange multiplier ($1/kT$). The expression of the density of probability has been discussed in Ref. [14]. It is shown in Ref. [15] that, for a canonical distribution of a system in contact with heat bath, the entropic index q is related to the particle number N and characterizes the bath. The Tsallis constant k is chosen as

$$\beta = \frac{Z_q^{1-q}}{c_q k_B T} \quad (13)$$

where k_B is the Boltzmann-Gibbs constant. The q -partition function Z_q is given by the normalisation of the probability density

$$Z_q = \int e_q(-\beta H) d\lambda \quad (14)$$

The measure μ defined relatively to the Lebesgue measure by $d\mu = (\rho^q/c_q) d\lambda$ is a measure of probability. In the extensive limit $q \rightarrow 1$, the q -Gibbs measure converges to the ordinary Boltzmann-Gibbs density. The substitution of the q -additivity Eq. (4) in the Tsallis entropy Eq. (1) gives

$$S_q = -k \int (-\beta H + \ln_q(Z_q^{-1}) - (1 - q)\beta H \ln_q(Z_q^{-1})) \rho^q d\lambda \quad (15)$$

and using Eq. (6)

$$S_q = k \int (\beta H + (1 - (1 - q)\beta H) Z_q^{q-1} \ln_q(Z_q)) \rho^q d\lambda \quad (16)$$

we find that the Tsallis entropy

$$S_q = k \int (\beta H \rho^q + \ln_q(Z_q) \rho) d\lambda \quad (17)$$

leads to the following formula

$$S_q = k\beta [H]_q + k \ln_q(Z_q)$$

Introducing the q -internal energy

$$U_q = \langle H \rangle_q \quad (18)$$

and the q -free energy

$$F_q = \frac{-1}{\beta} \ln_q(Z_q) \quad (19)$$

one finds

$$F_q = c_q U_q - T S_q \quad (20)$$

This relation is the q -generalisation of the ordinary equation of $S = U/T + k \ln(Z)$.

3 Mathematical Formalism

In this section, we introduce five important relations for the nonextensive mathematical formalism. The generalised internal energy U_q defined as the q -expectation of the Hamiltonian is given by

$$U_q = -\frac{1}{c_q} \frac{\partial \ln_q(Z_q)}{\partial \beta} \quad (21)$$

The proof of this equation is easy to compute.

$$\begin{aligned} -\frac{\partial \ln_q(Z_q)}{\partial \beta} &= -\frac{\partial \ln_q(Z_q)}{\partial Z_q} \frac{\partial Z_q}{\partial \beta} = \frac{-1}{Z_q^q} \frac{\partial Z_q}{\partial \beta} \quad (22) \\ &= \frac{-1}{Z_q^q} \frac{\partial}{\partial \beta} \int e_q(-\beta H) d\lambda \quad (23) \end{aligned}$$

Using the following expression of the derivative of the q -exponential function with respect to β

$$\frac{\partial e_q(-\beta H)}{\partial \beta} = -H e_q^q(-\beta H) \quad (24)$$

leads to

$$-\frac{\partial \ln_q(Z_q)}{\partial \beta} = \int H \frac{e_q^q(-\beta H)}{Z_q^q} d\lambda \quad (25)$$

$$= \int H \rho^q d\lambda = [H]_q = c_q U_q \quad (26)$$

Defining the q -movariance of two operators A and B by

$$Mov_q(A, B) = [AB]_{2q-1} - [A]_q [B]_q \quad (27)$$

the equation (Eq. 28) gives the q -movariance of A and H in function of the derivative of the q -mean of A

$$\frac{\partial [A]_q}{\partial \beta} - \left[\frac{\partial A}{\partial \beta} \right]_q = -q Z_q^{q-1} Mov_q(A, H) \quad (28)$$

The movariance converges to the usual covariance in the extensive limit $q \rightarrow 1$. Proof of the equation (Eq. 28) is

$$\begin{aligned} \frac{\partial [A]_q}{\partial \beta} &= \int \frac{\partial A}{\partial \beta} \rho^q d\lambda + \int A \frac{\partial \rho^q}{\partial \beta} d\lambda \quad (29) \\ &= \left[\frac{\partial A}{\partial \beta} \right]_q + q \int A \rho^{q-1} \frac{\partial}{\partial \beta} \left(\frac{e_q(-\beta H)}{Z_q} \right) d\lambda \quad (30) \end{aligned}$$

The derivative of the probability density with respect to the Lagrange multiplier is computed as

$$\begin{aligned} \frac{\partial \rho}{\partial \beta} &= \frac{1}{Z_q} \frac{\partial e_q(-\beta H)}{\partial \beta} - \frac{e_q(-\beta H)}{Z_q^2} \frac{\partial Z_q}{\partial \beta} \quad (31) \\ &= \frac{-H}{Z_q} e_q^q(-\beta H) + \frac{e_q(-\beta H)}{Z_q^2} \int H e_q^q(-\beta H) d\lambda \quad (32) \end{aligned}$$

Substituting the previous equation in Eq. (29) and using that Z_q is a constant leads to

$$\begin{aligned} \Delta_{q,\beta} A &= \frac{\partial [A]_q}{\partial \beta} - \left[\frac{\partial A}{\partial \beta} \right]_q \\ &= -q Z_q^{q-1} \int A H \rho^{2q-1} d\lambda + \dots \quad (33) \\ &\quad + q Z_q^{q-1} \int A \rho^q \left(\int H \rho^q d\lambda \right) d\lambda \quad (34) \\ &= -q Z_q^{q-1} ([AH]_{2q-1} - [A]_q [H]_q) \quad (35) \\ &= -q Z_q^{q-1} Mov_q(A, H) \quad (36) \end{aligned}$$

If the operator A does not depend explicitly of β , we find that the q -movariance is proportionnal to the derivative of the q -expectation of A with respect to β . We define the q -mariance by

$$Mar_q(A) = [A^2]_{2q-1} - [A]_q^2 \quad (37)$$

This function converges to the usual variance when q tends to 1. Using the equation of the generalised energy Eq. (21), we see that the q -mariance of the hamiltonian is given by

$$Mar_q(H) = \frac{1}{q Z_q^{q-1}} \frac{\partial^2 \ln_q(Z_q)}{\partial \beta^2} \quad (38)$$

Applying Cauchy-Schwarz inequality

$$\left(\int f g \, d\lambda \right)^2 \leq \left(\int f^2 \, d\lambda \right) \left(\int g^2 \, d\lambda \right) \quad (39)$$

with $f = A\rho^{q-1/2}Z_q^{q-1}$ and $g = \rho^{1/2}$ proves that $Mar_q(A)$ is always positive. In the extensive limit when q tends to 1, the variance of the hamitonian $Var(H) = \partial^2 \ln(Z)/\partial\beta^2$ is positive. Thus the internal energy as a function of the Lagrange multiplier β is a convex function.

We suppose now that the Hamiltonian H and the probability density function ρ depend on a parameter α distinct of β .

$$\frac{\partial \ln_q(Z_q)}{\partial \alpha} = \beta \left[\frac{\partial H}{\partial \alpha} \right]_q \quad (40)$$

Remark that this equation is also valid for α equal β (by Eq. (21)). Proof of this equation is easy to compute.

$$-\frac{\partial \ln_q(Z_q)}{\partial \alpha} = \frac{-1}{Z_q^q} \frac{\partial Z_q}{\partial \alpha} = \frac{-1}{Z_q^q} \frac{\partial}{\partial \alpha} \int e_q(-\beta H) \, d\lambda \quad (41)$$

Using the expression of the derivative of the q -exponential function with respect to α

$$\frac{\partial e_q(-\beta H)}{\partial \alpha} = -\beta \frac{\partial H}{\partial \alpha} e_q^q(-\beta H) \quad (42)$$

leads to

$$-\frac{\partial \ln_q(Z_q)}{\partial \beta} = \beta \int \frac{\partial H}{\partial \alpha} \rho^q \, d\lambda = \beta \left[\frac{\partial H}{\partial \alpha} \right]_q \quad (43)$$

The next fundamental equation gives the expression of the movariance of an observable and the derivation of the hamiltonian with respect to a given parameter

$$\frac{\partial [A]_q}{\partial \alpha} - \left[\frac{\partial A}{\partial \alpha} \right]_q = -q\beta Z_q^{q-1} Mov_q(A, \frac{\partial H}{\partial \alpha}) \quad (44)$$

The proof of this equation is

$$\begin{aligned} \frac{\partial [A]_q}{\partial \alpha} &= \int \frac{\partial A}{\partial \alpha} \rho^q \, d\lambda + \int A \frac{\partial \rho^q}{\partial \rho} \frac{\partial \rho}{\partial \alpha} \, d\lambda \\ &= \left[\frac{\partial A}{\partial \beta} \right]_q + q \int A \rho^{q-1} \frac{\partial}{\partial \alpha} \left(\frac{e_q(-\beta H)}{Z_q} \right) d\lambda \end{aligned} \quad (45)$$

The derivative of the partition function with respect to the parameter is given by

$$\frac{\partial Z_q}{\partial \alpha} = -\beta \int \frac{\partial H}{\partial \alpha} e_q^q(-\beta H) \, d\lambda \quad (47)$$

Let $\Delta_{q,\alpha}A$ be the quantity

$$\Delta_{q,\alpha}A = \frac{\partial [A]_q}{\partial \alpha} - \left[\frac{\partial A}{\partial \alpha} \right]_q \quad (48)$$

Substituing Eq. (42) and (47) in the previous equation and using that Z_q is a constant, we get

$$\Delta_{q,\alpha}A = q \int A \rho^{q-1} \frac{\partial}{\partial \alpha} \left(\frac{e_q(-\beta H)}{Z_q} \right) \, d\lambda \quad (49)$$

$$= -q Z_q^{q-1} \beta \int A \frac{\partial H}{\partial \alpha} \rho^{2q-1} \, d\lambda + \dots \quad (50)$$

$$+ q Z_q^{q-1} \beta \int A \rho^q \left(\int \frac{\partial H}{\partial \alpha} \rho^q \, d\lambda \right) \, d\lambda \quad (51)$$

$$= -q Z_q^{q-1} \left(\left[A \frac{\partial H}{\partial \alpha} \right]_{2q-1} - [A]_q \left[\frac{\partial H}{\partial \alpha} \right]_q \right) \quad (52)$$

$$= -q Z_q^{q-1} Mov_q(A, \frac{\partial H}{\partial \alpha}) \quad (53)$$

The following equation characterizes the derivation of the Tsallis entropy with respect to the α parameter

$$\frac{\partial S_q}{\partial \alpha} = -k\beta \left(\frac{\partial [H]_q}{\partial \alpha} - \left[\frac{\partial H}{\partial \alpha} \right]_q \right) \quad (54)$$

$$= qk\beta^2 Z_q^{q-1} Mov_q(H, \frac{\partial H}{\partial \alpha}) \quad (55)$$

In the extensive limit when q tends to 1, this equation shows that the derivation of the usual entropy is proportional to the variance of the hamiltonian

$$\frac{\partial S}{\partial \alpha} = \frac{1}{T} Var(H) = \frac{1}{T} \left(\frac{\partial \langle H \rangle}{\partial V} - \left\langle \frac{\partial H}{\partial V} \right\rangle \right) \quad (56)$$

If the parameter α is the volume V and the pressure defined by $P = -\langle \partial H / \partial V \rangle$, this equation is the usual thermodynamic relation $TdS = dU + PdV$.

If we define the q -pressure by

$$P_q = - \left\langle \frac{\partial H}{\partial V} \right\rangle_q = - \frac{1}{\beta c_q} \frac{\partial \ln_q(Z_q)}{\partial V} \quad (57)$$

and write the derivative of the Tsallis's entropy with respect to the volume

$$\frac{\partial S_q}{\partial V} = -k\beta c_q \left(\frac{\partial \langle H \rangle_q}{\partial V} - \left\langle \frac{\partial H}{\partial V} \right\rangle_q \right) \quad (58)$$

$$= -k\beta c_q \left(\frac{\partial U_q}{\partial V} + P_q \right) \quad (59)$$

leads to the ordinary equation

$$dU_q = -c_q T dS_q - P_q dV \quad (60)$$

Define the following q-quantities

$$\frac{\partial Q_q}{\partial V} = \frac{\partial \langle H \rangle_q}{\partial V} - \left\langle \frac{\partial H}{\partial V} \right\rangle_q \quad (61)$$

$$\frac{\partial W_q}{\partial V} = -P_q = \left\langle \frac{\partial H}{\partial V} \right\rangle_q \quad (62)$$

leads to the usual equation in the q-world

$$dU_q = dQ_q + dW_q = dQ_q - P_q dV \quad (63)$$

Derivating Eq. (8) with respect to parameter α

$$\frac{\partial S_q}{\partial \alpha} = \frac{k}{1-q} \frac{\partial c_q}{\partial \alpha} \quad (64)$$

Substituing the following equation

$$c_q = 1 + \frac{1-q}{k} S_q \quad (65)$$

and taking $A = 1$ in Eq. (44) leads to the differential equation follows by the Tsallis entropy.

$$\begin{aligned} & \frac{\partial S_q}{\partial \alpha} - q\beta Z_q^{q-1} \frac{\partial [H]_q}{\partial \alpha} S_q \\ &= \frac{q}{q-1} k\beta Z_q^{q-1} \left(\frac{\partial [H]_{2q-1}}{\partial \alpha} - \frac{\partial [H]_q}{\partial \alpha} \right) \end{aligned} \quad (66)$$

4 Ideal Gas

The classical ideal gas in D-dimensional space has been exhaustively discussed in Ref. [16]. Correlations were analyzed in Ref.[17]. We summarize some results in the case $0 < q < 1$ and show by straightforward calculation how the partition function is computed. The hamiltonian is

$$H = \sum_{i=1}^N \frac{p_i^2}{2m} \quad (68)$$

where m is the particle mass, N the particle number and p_i the momentum of the i th particle. The partition function is given by

$$Z_q = \int_{\Omega} (1 - a(p_1^2 + \dots + p_N^2))^{\frac{1}{1-q}} \frac{d^D p_i d^D r_i}{N! h^{DN}} \quad (69)$$

where a is the positive constant $a = (1-q)\beta/2m$ and Ω is domain of integration such that $1 - a(p_1^2 + \dots + p_N^2) > 0$.

$$Z_q = \frac{V^N}{N! h^{DN}} \int_{\Omega} (1 - a(p_1^2 + \dots + p_N^2))^{\frac{1}{1-q}} d^D p_i \quad (70)$$

and

$$Z_q = \frac{V^N}{N! h^{DN}} \frac{2\pi^{DN/2}}{\Gamma(DN/2)} \int_0^{1/\sqrt{a}} (1 - ar^2)^{\frac{1}{1-q}} r^{\frac{DN}{2}-1} dr \quad (71)$$

Let x be ar^2 . Using the fact that the beta distribution is a density of probability for two parameters u and v greatest or equal 1, that is

$$\frac{\Gamma(u+v)}{\Gamma(u)\Gamma(v)} \int_0^1 x^{u-1} (1-x)^{v-1} dx = 1 \quad (72)$$

After a simple calculation, we find the following expression of the q -partition function

$$Z_q = \frac{V^N}{N!} \left(\frac{2\pi m}{(1-q)\beta h^2} \right)^{DN/2} \frac{\Gamma(\frac{2-q}{1-q})}{\Gamma(\frac{2-q}{1-q} + \frac{DN}{2})} \quad (73)$$

The same calculation for the constant c_q leads to

$$c_q = Z_q^{1-q} \frac{\Gamma(\frac{1}{1-q})}{\Gamma(\frac{1}{1-q} + \frac{DN}{2})} \frac{\Gamma(\frac{2-q}{1-q} + \frac{DN}{2})}{\Gamma(\frac{2-q}{1-q})} \quad (74)$$

Using the following property of the gamma function

$$\Gamma(z+1) = z\Gamma(z) \quad (75)$$

we find

$$c_q = Z_q^{1-q} (1 + (1-q) \frac{DN}{2}) \quad (76)$$

The generalised energy given by Eq. (21) is computed in the same way

$$U_q = \frac{DN}{2\beta} \frac{Z_q^{1-q}}{c_q} \quad (77)$$

Using the expression of the Lagrange multiplier given by Eq. (29), we obtain the following expression of the internal energy which is independent of the entropic index.

$$U_q = \frac{DN}{2} k_B T \quad (78)$$

The Tsallis and Boltzmann-Gibbs constants are related to the equation

$$k = c_q Z_q^{q-1} k_B = \left(1 - (1-q) \frac{DN}{2}\right) k_B \quad (79)$$

The equation of state

$$P_q = \frac{1}{\beta c_q Z_q^q} \frac{\partial Z_q}{\partial V} = \frac{N}{V} \frac{Z_q^{1-q}}{\beta c_q} \quad (80)$$

leads to the usual expression

$$P_q V = N k_B T \quad (81)$$

In the extensive limit when q tends to 1, we recover the equation of ideal gas. Remark that if we replace the hamiltonian by the same expression minus the generalized energy, the results are unchanged [18]. The q -mariance of the hamitonian

$$Marq(H) = \frac{1}{q} k_B T U_q \quad (82)$$

converges in the extensive limit to the variance of the hamiltonian $DN k_B^2 T^2 / 2$. Simple calculations give the movariance of the q -pressure

$$Mov_q(H, P_q) = \frac{1-q}{q} \frac{DN}{2\beta} P_q c_q Z_q^{1-q} \quad (83)$$

and

$$Mov_q(H, \frac{\partial H}{\partial V}) = \frac{-Nk}{q V k_B \beta^2} Z_q^{2-2q} \quad (84)$$

and the ratio

$$\frac{Mov_q(H, P_q)}{Mov_q(H, \frac{\partial H}{\partial V})} = (q-1) \frac{k}{k_B} \frac{DN}{2} \quad (85)$$

5 Quantum Free Particle

In this section, a free particle confined in a 3-dimensional box is considered. Denoting L_1 , L_2 ,

L_3 the sides of the box, it is well known that the energy eigenvalues are given by

$$E = \frac{h^2}{8m} \left(\frac{n_1^2}{L_1^2} + \frac{n_2^2}{L_2^2} + \frac{n_3^2}{L_3^2} \right) \quad (86)$$

where the quantum numbers are non-negative integers $n_i = 0, 1, 2, \dots$. The partition function is given by

$$Z_q = \sum_{n_1, n_2, n_3} e_q(-\beta E) = \sum_{n_1, n_2, n_3} (1 - (1-q)E)^{\frac{1}{1-q}} \quad (87)$$

Let λ_T be the translation temperature

$$\lambda_T = \frac{h^2}{8mkL^2} \quad (88)$$

where $L \sim L_i$ is the length of a side. For sufficiently large $T \gg \lambda_T$, the sum over n_i can be replaced by an integral

$$Z_q = \frac{1}{8} \int (1 - (1-q)\beta E)^{\frac{1}{1-q}} 1_\Omega(n_i) dn_1 dn_2 dn_3 \quad (89)$$

where $\Omega = \{1 - (1-q)\beta E > 0\}$ and $1_A(x) = 1$ if $x \in A$ and 0 otherwise. For a volume $V = L_1 L_2 L_3$ and a parameter $b = \beta(1-q)h^2/8m$ the change of variables

$$Z_q = \frac{V}{8} \frac{2\pi^{3/2}}{\Gamma(3/2)} \int_0^{1/\sqrt{b}} (1 - br^2)^{\frac{1}{1-q}} r^2 dr \quad (90)$$

could be written as a beta integral

$$Z_q = \frac{V}{8} \frac{\pi^{3/2}}{b^{3/2} \Gamma(3/2)} \int_0^1 (1-x)^{\frac{1}{1-q}} x^{\frac{1}{2}} dx \quad (91)$$

thus

$$Z_q = V \left(\frac{2\pi}{\beta(1-q)h^2} \right)^{3/2} \frac{\Gamma(\frac{2-q}{1-q})}{\Gamma(\frac{2-q}{1-q} + \frac{3}{2})} \quad (92)$$

this is the classical formula with $N = 1$ and $D = 3$.

6 Fermions and Bosons Distributions

Thermodynamics on quantum groups and thermodynamics of boson and fermion systems with fractal

distributions have been studied recently [19, 20, 21]. In the grand canonical ensemble, the grand partition function is the sum under the states j

$$\Theta_q = \sum_j (1 - \beta(1 - q)(E_j - \mu))^{\frac{1}{1-q}} \quad (93)$$

where μ is the chemical potential. The density of states in the phase space is given by

$$D(E) = \frac{\partial}{\partial E} \int \frac{d^3r d^3p}{h^3} = \frac{2\pi V}{h^3} (2m)^{3/2} \sqrt{E} \quad (94)$$

In the thermodynamic limit, the sum is replaced by a continuous integration, thus the average is defined by

$$\langle A \rangle_q = \sum_j A_j p_j^q \simeq \int_0^\infty A D(E) f^q(E) dE \quad (95)$$

where f is the function

$$f(E) = \frac{1}{(1 - \beta(1 - q)(E - \mu))^{\frac{1}{1-q}}} 1_\Omega \quad (96)$$

The change of variable $E = x^2/\beta(q - 1)$ gives

$$\langle A \rangle_q = \frac{4\pi V}{h^3} \left(\frac{2mkT}{q - 1} \right)^{3/2} \int_0^\infty A^q x^2 f(x) dx \quad (97)$$

The average total number of particles

$$\bar{N} = \sum_j \frac{1}{(1 - \beta(1 - q)(E_j - \mu))^{\frac{1}{1-q}}} \quad (98)$$

in the thermodynamic limit

$$\bar{N} \simeq \int_0^\infty D(E) f(E) dE \quad (99)$$

is calculated by

$$\bar{N} = K \int_a^\infty \frac{x^2}{(1 + x^2 - \beta(q - 1)\mu)^{\frac{1}{1-q}}} dx \quad (100)$$

where $a^2 = -1 + \beta(q - 1)\mu$ and

$$K = \frac{4\pi V}{h^3} \left(\frac{2mkT}{q - 1} \right)^{3/2} \quad (101)$$

This integral could be computed as a beta integral if $1 \leq q \leq 7/5$

$$\bar{N} = \frac{4\pi V}{3(q - 1)h^3} \left(\frac{2mkT}{q - 1} \right)^{3/2} a^{-2u} \frac{\Gamma(u)\Gamma(v)}{\Gamma(u + v)} \quad (102)$$

where $u = (5 - 3q)/2(q - 1)$ and $v = 1/(1 - q)$. Same calculation could be done for the generalised energy in the thermodynamic limit

$$U_q = \frac{2\pi V}{h^3} \int_0^\infty (2mE)^{3/2} f^q(E) dE \quad (103)$$

after simple manipulation

$$U_q = \frac{4\pi V}{2\beta(q - 1)h^3} \left(\frac{2mkT}{q - 1} \right)^{3/2} a^{-2u} \frac{\Gamma(u)\Gamma(v)}{\Gamma(u + v)} \quad (104)$$

The average energy per particle

$$U_q = \frac{3}{2} \bar{N} kT \quad (105)$$

It is worth noting that the generalized energy is in this equation related to the Tsallis constant (and not to the Boltzmann-Gibbs constant).

7 Conclusions

In this paper, we studied the probabilistic properties within the framework of Tsallis thermodynamics. It has been shown that movariance and marivariance are suitable functions for generalized covariance and variance in the non-extensive thermodynamic context. The invariance of thermodynamic relations is established if the Tsallis constant depend on the entropic parameter. However, in the canonical ensemble, as illustrated on the ideal gas, the equation of state has the same form as the ordinary equation in the non-extensive formalism with the Boltzmann-Gibbs constant. In the grand canonical ensemble, the dependence of the q -average internal energy is related to the Tsallis constant. In both cases, we recover the usual equations in the extensive limit.

Acknowledgements

I would like to thank Professor Tsallis for making valuable suggestions to improve this paper and A.K. Rajagopal for fruitful conversations and perceptive comments.

References

- [1] S. Abe, S. Martinez, F. Pennini and A. Plastino, Phys. Lett. A (2001), in press [cond-mat/0011012], (2001).
- [2] C. Tsallis, J. Stat. Phys. **52**, 479 (1988).
- [3] C. Tsallis, Braz. J. Phys. **29**, 1 (1999). This is a special issue on Tsallis thermostatics and can be obtained from http://www.sbf.if.usp.br/WWW_pages/Journals/BJP/Vol129/Num1/index.htm. See also <http://tsallis.cat.cbpf.br/biblio.htm> for a regularly updated bibliography on the subject.
- [4] D.H. Zanette and P. A. Alemany, Phys. Rev. Lett. **75**, 366 (1995).
- [5] J. Naudts, Dual description of nonextensive ensembles, to appear in the Proc. of the “*International Workshop on Classical and Quantum Complexity and Nonextensive Thermodynamics*” (Denton, Texas, 3-6 April 2000), eds. P. Grigolini, C. Tsallis and B.J. West, *Chaos, Solitons and Fractals* (2001), in press.[cond-mat/9904070].
- [6] S. Abe and A. K. Rajagopal, J. Phys. A **33**, 8733, (2000), preprint [cond-mat/0002159].
- [7] S. Abe, S. Martinez, F. Pennini, A. Plastino, Phys. Lett. A (2001), in press [cond-mat/0006109], (2000).
- [8] A. K. Rajagopal, Phys. Rev. Lett. **76**, 3469 (1996).
- [9] A. K. Rajagopal, Braz. J. Phys **29**, 61 (1999).
- [10] S. Abe, Y. Okamoto (eds), *Nonextensive Statistical Mechanics and its Applications*, Springer (2001).
- [11] C. Tsallis, Phys. Rev. Lett. **75**, 3589 (1995).
- [12] F. Pennini, A. R. Plastino and A. Plastino, Physica A **258**, 446 (1998).
- [13] W. Gibbs, *Elementary Principles in Statistical Mechanics* (Yale University Press, New Haven, 1902).
- [14] D. Prato and C. Tsallis, Phys. Rev. E **60**, 2398 (1999).
- [15] A. R. Plastino and A. Plastino, Phys. Lett. A **193**, 140 (1994).
- [16] S. Abe, Phys. Lett. A **263** (1999) 424; Erratum **267** (2000) 456.
- [17] S. Abe, Physica A **269** (1999) 403.
- [18] A. K. Rajagopal and S. Abe, Phys. Rev. Lett. **83**, 1711 (1999).
- [19] M.R. Ubriaco, Phys. Rev. E **57**, 179 (1998).
- [20] M.R. Ubriaco, Phys. Rev. E **60**, 165 (1999).
- [21] E.K. Lenzi, R.S. Mendel and A. K. Rajagopal, Physica A **286**, 503, [cond-mat/9904100], (2000).
- [22] R. Balian, N.L. Balazs, Ann. Phys., (NY) **179**, 97 (1987).